


Tbp-Flox Genotyping Protocol




<i>Common Name</i>	<i>Tbp-Flox</i>	<i>Cat. NO.</i>	<i>NM-CKO-240133</i>
<i>Strain of Origin</i>	<i>C57BL/6J</i>	<i>Version</i>	<i>V1</i>

Genotyping strategy

Wild type allele



Targeted allele




→ : primer location

Primers

<i>Primer</i>	<i>Sequence (5'→3')</i>	<i>Primer type</i>
<i>P1</i>	<i>GACACGTGCTTAGTCAGTGGA</i>	<i>Forward</i>
<i>P2</i>	<i>AGATGTTTAGCCTCTCATGTGT</i>	<i>Reverse</i>

Expected results

Results	
Genotype	Wild type: P1P2 =270 bp; Heterozygote: P1P2 =329 bp and 270 bp; Homozygote: P1P2 =329 bp

Reaction &Cycling

PCR Reaction System	Reaction Component			Volume (μl)
	ddH ₂ O			8.0
	2 × Rapid Taq Master Mix			10.0
	P1(10 pmol/μl)			0.5
	P2(10 pmol/μl)			0.5
	Genomic DNA			1.0
	Total			20
	2 × Rapid Taq Master Mix from Vazyme(Code Number: P222-01)			
Cycling	Step	Temp	Time	Note
Reaction	1	95°C	5 min	

	2	95°C	20 sec	
	3	60°C	20 sec	
	4	72°C	20 sec	35 repeats to 2
	5	72°C	5 min	
	6	12°C	Hold	