


Bicc1-Flox Genotyping Protocol




Common Name	Bicc1-Flox	Cat. NO.	NM-CKO-240848
Strain of Origin	C57BL/6J	Version	V1

Genotyping strategy

Wild type allele



Targeted allele




→ : primer location

Primers

Primer	Sequence (5'→3')	Primer type
P1	TGCACTATAGGGCCAGGTCT	Forward
P2	GCAAATCACACCACTGCGAA	Reverse

Expected results



Genotype	Wild type: P1P2 =234 bp; Heterozygote: P1P2 =293 bp and 234 bp; Homozygote: P1P2 =293 bp
----------	--

Reaction &Cycling

PCR Reaction System	Reaction Component			Volume (μl)
	ddH ₂ O			8.0
	2 × Rapid Taq Master Mix			10.0
	P1(10 pmol/μl)			0.5
	P2(10 pmol/μl)			0.5
	Genomic DNA			1.0
	Total			20
	2 × Rapid Taq Master Mix from Vazyme(Code Number: P222-01)			
Cycling Reaction	Step	Temp	Time	Note
	1	95°C	5 min	
	2	95°C	20 sec	
	3	60°C	20 sec	
	4	72°C	20 sec	35 repeats to 2
	5	72°C	5 min	
	6	12°C	Hold	