


Cldn24-KO Genotyping Protocol



Common Name	Cldn24-KO	Cat. NO.	NM-KO-241914
Strain of Origin	C57BL/6J	Version	V1


Genotyping strategy



Primers

Primer	Sequence (5'→3')	Primer type
P1	ACGTTAGGGTGAGGGGTCAT	Forward
P2	AATCCCCAGCATACACAGGC	Reverse
P3	ACTCGTCAAGCACTGGGGAT	Forward
P4	TGGATGACGCAGGATTTCCA	Reverse

Expected results

Results	 GeneRuler™ 1 kb DNA Ladder 1% Tris/borate/EDTA agarose (pH 8.0) 0.5 µg/lane, 8 cm length gel, 1X TAE, 7 V/cm, 45 min DL2000 bp 2000 1000 750 500 250 100 1.7% agarose
Genotype	Knockout type: -759bp Wild type: P1P2 =1261 bp; P3P4 =446 bp Heterozygote: P1P2 =1261 bp and 502 bp; P3P4 =446 bp Homozygote: P1P2 =502 bp

Note: In both wild-type and heterozygous mice, whether the P1 and P2 primers can amplify larger bands does not affect the interpretation of the results, because the purpose of designing this pair of primers is to amplify KO band

Reaction & Cycling

PCR Reaction System	Reaction Component			Volume (µl)
	ddH ₂ O			8.0
	2×Rapid Taq Master Mix			10.0
	P1(10 pmol/µl) or P3(10 pmol/µl)			0.5
	P2(10 pmol/µl) or P4(10 pmol/µl)			0.5
	Genomic DNA			1.0
	Total			20
	2×Rapid Taq Master Mix from Vazyme(Code Number: P222-01)			
Cycling Reaction	Step	Temp	Time	Note
	1	95°C	5 min	
	2	95°C	20 sec	
	3	60°C	20 sec	
	4	72°C	20 sec	35 repeats to 2
	5	72°C	5 min	
	6	12°C	Hold	